

On the Time-Modulation of the β^+ -Decay Rate of H-like $^{140}\text{Pr}^{58+}$ Ion

A. N. Ivanov^{a,b}, E. L. Kryshen^c, M. Pitschmann^a, P. Kienle^{b,d}

^a *Atominstitut der Österreichischen Universitäten, Technische Universität Wien,
Wiedner Hauptstrasse 8-10, A-1040 Wien, Österreich*

^b *Stefan Meyer Institut für subatomare Physik Österreichische Akademie der Wissenschaften,
Boltzmannngasse 3, A-1090, Wien, Österreich*

^c *Petersburg Nuclear Physics Institute, 188300 Gatchina, Orlova roscha 1, Russian Federation and*

^d *Physik Department, Technische Universität München, D-85748 Garching, Germany**

(Dated: June 16, 2008)

According to recent experimental data at GSI, the rates of the number of daughter ions, produced by the nuclear K-shell electron capture (EC) decays of the H-like ions $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$, are modulated in time with periods $T_{EC} \simeq 7$ sec and amplitudes $a_{EC} \simeq 0.20$. We study a possible time-dependence of the nuclear positron (β^+) decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion. We show that the time-dependence of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion as well as any H-like heavy ions cannot be observed. This result can be used as a prediction for future analysis of the time-dependence of the β^+ -decay rates of the H-like heavy ions $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$ at GSI for the test of the measuring method.

PACS: 12.15.Ff, 13.15.+g, 23.40.Bw, 26.65.+t

INTRODUCTION

The experimental investigation of the EC-decays of the H-like ions $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$, $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{58+} + \nu$ and $^{142}\text{Pm}^{60+} \rightarrow ^{142}\text{Nd}^{60+} + \nu$, carried out at the Experimental Storage Ring (ESR) at GSI in Darmstadt [1], showed a modulation in time with periods $T_{EC} \simeq 7$ s of the rates of the number $N_d^{EC}(t)$ of daughter ions $^{140}\text{Ce}^{58+}$ and $^{142}\text{Nd}^{60+}$, respectively (see Fig. 1). Since the rate of the number of daughter ions is defined by [1]

$$\frac{dN_d^{EC}(t)}{dt} = \lambda_{EC}(t) N_m(t), \quad (1)$$

where $N_m(t)$ is the number of mother $^{140}\text{Pr}^{58+}$ or $^{142}\text{Pm}^{60+}$ ions, a periodic time-dependence can be attributed to the time-dependent EC-decay rate $\lambda_{EC}(t)$ [1]

$$\lambda_{EC}(t) = \lambda_{EC} \left\{ 1 + a_{EC} \cos \left(\frac{2\pi t}{T_{EC}} + \phi \right) \right\}, \quad (2)$$

where ϕ and a_{EC} are the phase and the amplitude of the time-dependent term, which are not well-measured quantities [2, 3].

In turn, periods of the time-modulation T_{EC} are measured well and equal to $T_{EC} = 7.06(8)$ s and $T_{EC} = 7.10(22)$ s for $^{140}\text{Ce}^{58+}$ and $^{142}\text{Nd}^{60+}$, respectively [1]. Such a periodic dependence has been explained in [4, 5, 6] in terms of the massive neutrino mixing with the period T_{EC} equal to

$$T_{EC} = \frac{4\pi\gamma M_m}{(\Delta m_{21}^2)_{\text{GSI}}}, \quad (3)$$

where M_m is the mass of the mother ion [6], $\gamma = 1.43$ is the Lorentz factor [1] and $(\Delta m_{21}^2)_{\text{GSI}} =$

$2.22(4) \times 10^{-4} \text{ eV}^2$, calculated in [4] for the experimental value $T_{EC} = 7$ s [1]. A relation of the value $(\Delta m_{21}^2)_{\text{GSI}} = 2.22(4) \times 10^{-4} \text{ eV}^2$ to the KamLAND experimental data $(\Delta m_{21}^2)_{\text{KamLAND}} = 0.80^{+0.06}_{-0.05} \times 10^{-4} \text{ eV}^2$ [7] has been investigated and obtained in [5].

The EC-decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion, averaged over time $\langle \lambda_{EC}(t) \rangle = \lambda_{EC}$, as well as the β^+ -decay rate of the $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{57+} + e^+ + \nu$ decay have been also measured at GSI [8]: $\lambda_{EC}^{\text{exp}} = 0.00219(6) \text{ s}^{-1}$ and $\lambda_{\beta^+}^{\text{exp}} = 0.00161(10) \text{ s}^{-1}$ with the ratio $R_{EC/\beta^+}^{\text{exp}} = 1.36(9)$.

The calculation of the EC and β^+ decay rates of the H-like $^{140}\text{Pr}^{58+}$ ion as well as the He-like $^{140}\text{Pr}^{57+}$ ion has been carried out in [9]

$$\begin{aligned} \lambda_{EC} &= \frac{1}{2F+1} \frac{3}{2} |\mathcal{M}_{\text{GT}}|^2 |\langle \psi_{1s}^{(Z)} \rangle|^2 \frac{Q_H^2}{\pi}, \\ \lambda_{\beta^+} &= \frac{2}{2F+1} \frac{|\mathcal{M}_{\text{GT}}|^2}{4\pi^3} f(Q_{\beta^+}, Z-1), \end{aligned} \quad (4)$$

where $F = 1/2$ is the total angular momentum of the H-like $^{140}\text{Pr}^{58+}$, $Q_H = (3348 \pm 6) \text{ keV}$ and $Q_{\beta^+} = (3396 \pm 6) \text{ keV}$ are the Q -values of the $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{58+} + \nu$ and $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{57+} + e^+ + \nu$ decays, respectively; $\mathcal{M}_{\text{GT}}$ is the nuclear matrix element of the Gamow-Teller transition

$$\mathcal{M}_{\text{GT}} = -2g_A G_F V_{ud} \int d^3x \Psi_d^*(r) \Psi_m(r), \quad (5)$$

where $\Psi_d^*(r)$ and $\Psi_m(r)$ are the wave functions of the daughter and mother nuclei, respectively, and $\Psi_d^*(r) \Psi_m(r) \sim \rho(r)$ is the nuclear matter density $\rho(r)$ having the Woods-Saxon shape with $R = 1.1 A^{1/3} \text{ fm}$ and diffuseness parameter $a = 0.50 \text{ fm}$ [10]. Then, $\psi_{1s}^{(Z)}$ is the Dirac wave function of




FIG. 1: The experimental non-exponential time-dependence of the rate of the number of daughter ions, given by Eq.(1), in the EC -decays $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{58+} + \nu$ and $^{142}\text{Pm}^{60+} \rightarrow ^{142}\text{Nd}^{60+} + \nu$ of the H-like $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$ ions, respectively.

the bound electron in the ground state, $Z = 59$ is the electric charge of the mother nucleus $^{140}\text{Pr}^{59+}$. The average value of the Dirac wave function of the bound electron $\langle \psi_{1s}^{(Z)} \rangle$ is defined by [9]

$$\langle \psi_{1s}^{(Z)} \rangle = \frac{\int d^3x \psi_{1s}^{(Z)}(\vec{r}) \rho(r)}{\int d^3x \rho(r)} = \frac{1.66}{\sqrt{\pi a_B^3}} \quad (6)$$

where $a_B = 1/m_e Z \alpha = 897 \text{ fm}$ for the electron mass $m_e = 0.511 \text{ MeV}$ and the fine-structure constant $\alpha = 1/137.036$. In the β^+ -decay rate $f(Q_{\beta^+}, Z-1) = (2.21 \pm 0.03) \text{ MeV}^5$ is the Fermi integral [9]. The theoretical prediction for the ratio $R_{EC/\beta^+}^{\text{th}}$ is [9]

$$R_{EC/\beta^+}^{\text{th}} = \frac{3\pi^2 Q_H^2 |\langle \psi_{1s}^{(Z)} \rangle|^2}{f(Q_{\beta^+}, Z-1)} = 1.40(4), \quad (7)$$

which agrees well with the experimental data $R_{EC/\beta^+}^{\text{exp}} = 1.36(9)$.

According to [2, 3], a time-dependence of the rate of the number of daughter ions in the β^+ -decay of the H-like $^{140}\text{Pr}^{58+}$ ion has not been studied experimentally until now.

In this paper we apply a theoretical approach, developed in [4]–[6] for the analysis of the time-modulation in the EC -decay of the H-like $^{140}\text{Pr}^{58+}$, to the study of the time-dependence

of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion. Its experimental investigation should be a stringent test of the applied single-ion Schottky mass-measurement method.

AMPLITUDES OF THE β^+ -DECAY OF THE H-LIKE $^{140}\text{Pr}^{58+}$ ION

Following [4], for the calculation of the time-modulation of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion we use time-dependent perturbation theory. The Hamilton operator $H_W(t)$ of the weak interactions is given by

$$\begin{aligned} H_W(t) &= \frac{G_F}{\sqrt{2}} V_{ud} \sum_j U_{ej} \\ &\times \int d^3x [\bar{\psi}_n(x) \gamma^\mu (1 - g_A \gamma^5) \psi_p(x)] \\ &\times [\bar{\psi}_{\nu_j}(x) \gamma_\mu (1 - \gamma^5) \psi_e(x)] \end{aligned} \quad (8)$$

with standard notations [4]. In our analysis neutrinos ν_j ($j = 1, 2, 3$) are Dirac particles with masses m_j ($j = 1, 2, 3$) [7].

The amplitude of the β^+ -decay of the H-like $^{140}\text{Pr}^{58+}$ ion with undetected neutrino we define as a coherent sum of the amplitudes of the β^+ -decays $^{140}\text{Pr}^{58+} \rightarrow ^{140}\text{Ce}^{57+} + e^+ + \nu_j$ [4, 6]

$$\begin{aligned} \mathcal{M}_{FM_F \rightarrow F' M_{F'}}(t) &= -i \sum_j \int_{-\infty}^t d\tau \\ &\times \langle \nu_j(\vec{k}_j) e^+(p_+) d(\vec{q}) | H_W(\tau) | m(\vec{0}) \rangle d\tau, \end{aligned} \quad (9)$$

where m and d are the mother ion $^{140}\text{Pr}^{58+}$ and the daughter ion $^{140}\text{Ce}^{57+}$, respectively. The mother ion is taken in the state $^{140}\text{Pr}_{F=\frac{1}{2}}^{58+}$ with $F = \frac{1}{2}$ and $M_F = \pm \frac{1}{2}$ and in the rest frame [4]. In turn, the daughter $^{140}\text{Ce}^{57+}$ ion is a H-like ion in the state with $F' = \frac{1}{2}$ and $M_{F'} = \pm \frac{1}{2}$.

The wave function of the neutrino ν_j we define in the form of a wave packet [4, 6]

$$\begin{aligned} \psi_{\nu_j}(\vec{r}, t) &= (2\pi\delta^2)^{3/2} \int \frac{d^3k}{(2\pi)^3} e^{-\frac{1}{2}\delta^2(\vec{k} - \vec{k}_j)^2} \\ &\times e^{i\vec{k} \cdot \vec{r} - iE_j(\vec{k})t} u_{\nu_j}(\vec{k}, \sigma_j). \end{aligned} \quad (10)$$

where a spatial smearing of the neutrino ν_j is determined by the parameter δ , $\vec{k}_j$ is the neutrino momentum and $E_j(\vec{k}) = \sqrt{\vec{k}^2 + m_j^2}$ is the energy of a plane wave with the momentum $\vec{k}$, $u_{\nu_j}(\vec{k}, \sigma_{\nu_j})$ is the Dirac bispinor of the neutrino ν_j [4]. In the limit $\delta \rightarrow \infty$, due to the relation $(2\pi\delta^2)^{3/2} e^{-\frac{1}{2}\delta^2(\vec{k} - \vec{k}_j)^2} \rightarrow (2\pi)^3 \delta^{(3)}(\vec{k} - \vec{k}_j)$,

the wave function (10) reduces to the form of a plane wave [4].

Following [4] and [9] we obtain the amplitudes of the β^+ -decay

$$\begin{aligned}
\mathcal{M}_{\frac{1}{2}, +\frac{1}{2} \rightarrow \frac{1}{2}, M'_F}(t) &= -\sqrt{2M_m 2E_d} \frac{\mathcal{M}_{\text{GT}}}{2\sqrt{2}} (\pi\delta^2)^{3/2} \\
&\times e^{\varepsilon t} \sum_j U_{ej} e^{-\frac{1}{2}(\vec{k}_d + \vec{p}_+ + \vec{k}_j)^2} \frac{e^{i\Delta E_j(\vec{k}_j)t}}{\Delta E_j(\vec{k}_j) - i\varepsilon} \\
&\times \left\{ \sqrt{\frac{2}{3}} [\varphi_{n, -\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, +\frac{1}{2}}] \cdot [\bar{u}_{\nu_j}(\vec{k}_j, \sigma_j) \vec{\gamma} (1 - \gamma^5)] \right. \\
&\times v_{e^+}(\vec{p}_+, \sigma_+) \delta_{M'_F, -\frac{1}{2}} - \sqrt{\frac{1}{6}} \\
&\times [\varphi_{n, +\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, +\frac{1}{2}} - \varphi_{n, -\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, -\frac{1}{2}}] \cdot [\bar{u}_{\nu}(\vec{k}_j, \sigma_j) \vec{\gamma}] \\
&\times (1 - \gamma^5) v_{e^+}(\vec{p}_+, \sigma_+) \delta_{M'_F, +\frac{1}{2}} \left. \right\}, \\
\mathcal{M}_{\frac{1}{2}, -\frac{1}{2} \rightarrow \frac{1}{2}, M'_F}(t) &= -\sqrt{2M_m 2E_d} \frac{\mathcal{M}_{\text{GT}}}{2\sqrt{2}} (\pi\delta^2)^{3/2} \\
&\times e^{\varepsilon t} \sum_j U_{ej} e^{-\frac{1}{2}(\vec{k}_d + \vec{p}_+ + \vec{k}_j)^2} \frac{e^{i\Delta E_j(\vec{k}_j)t}}{\Delta E_j(\vec{k}_j) - i\varepsilon} \\
&\times \left\{ \sqrt{\frac{2}{3}} [\varphi_{n, +\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, -\frac{1}{2}}] \cdot [\bar{u}_{\nu}(\vec{k}_j, \sigma_j) \vec{\gamma} (1 - \gamma^5)] \right. \\
&\times v_{e^+}(\vec{p}_+, \sigma_+) \delta_{M'_F, +\frac{1}{2}} - \sqrt{\frac{1}{6}} \\
&\times [\varphi_{n, +\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, +\frac{1}{2}} - \varphi_{n, -\frac{1}{2}}^\dagger \vec{\sigma} \varphi_{p, -\frac{1}{2}}] \cdot [\bar{u}_{\nu}(\vec{k}_j, \sigma_j) \vec{\gamma}] \\
&\times (1 - \gamma^5) v_{e^+}(\vec{p}_+, \sigma_+) \delta_{M'_F, -\frac{1}{2}} \left. \right\}. \quad (11)
\end{aligned}$$

For the calculation of the amplitudes Eq.(11) we have carried out the integration over $\vec{k}$ with the δ -function $(2\pi)^3 \delta^{(3)}(\vec{k} + \vec{k}_d + \vec{p}_+)$ and denoted $\Delta E_j(\vec{k}_j) = E_d(\vec{k}_d) + E_+(\vec{p}_+) + E_j(\vec{k}_j) - M_m$, where $E_d(\vec{k}_d)$, $E_+(\vec{p}_+)$ and $E_j(\vec{k}_j)$ are energies of the daughter ion, positron and neutrino ν_j , $\vec{k}_d$, $\vec{p}_+$ and $\vec{k}_j$ are their momenta and M_m is the mother ion mass. For the calculation of the integral over time we have used the ε -regularization [4]. Finally the parameter ε should be taken in the limit $\varepsilon \rightarrow 0$.

TIME-DEPENDENCE OF THE β^+ -DECAY RATE OF THE H-LIKE $^{140}\text{Pr}^{58+}$ ION

According to [4], the first step to the calculation of the time-dependent β^+ -decay rate $\lambda_{\beta^+}^{(\text{H})}(t)$ of the H-like $^{140}\text{Pr}^{58+}$ ion is the calculation of the rate of the neutrino spectrum $N_\nu(t)$. It is defined by [4]

$$\frac{dN_\nu(t)}{dt} =$$

$$\begin{aligned}
&= \frac{1}{2M_m} \frac{1}{2F+1} \frac{d}{dt} \int \sum_{M_F, M'_F} |\mathcal{M}_{F, M_F \rightarrow F', M'_F}(t)|^2 \\
&\times F(Z-1, E_+) \frac{d^3 k_d}{(2\pi)^3 2E_d} \frac{d^3 p_+}{(2\pi)^3 2E_+} \quad (12)
\end{aligned}$$

where $F(Z-1, E_+)$ is the Fermi function [11] (see also [9]), describing the Coulomb repulsion between the positron and the nucleus $^{140}\text{Ce}^{58+}$. It is equal to [11]

$$\begin{aligned}
F(Z-1, E_+) &= \\
&= \left(1 + \frac{1}{2}\gamma\right) \frac{4(2Rp_+)^{2\gamma}}{\Gamma^2(3+2\gamma)} e^{-\frac{\pi(Z-1)\alpha E_+}{p_+}} \\
&\times \left| \Gamma\left(1 + \gamma - i \frac{\alpha(Z-1)E_+}{p_+}\right) \right|^2, \quad (13)
\end{aligned}$$

where $p_+ = \sqrt{E_+^2 - m_e^2}$, $R = 5.712 \text{ fm}$, $Z = 59$ and $\gamma = \sqrt{1 - ((Z-1)\alpha)^2} - 1$.

After the integration over the phase volume of the daughter ion, the directions of the positron momentum $\vec{p}_+$ and the limit $\varepsilon \rightarrow 0$ we arrive at the following expression for the rate of the neutrino spectrum

$$\begin{aligned}
\frac{dN_\nu(t)}{dt} &= \frac{1}{2F+1} \frac{|\mathcal{M}_{\text{GT}}|^2}{\pi^2} (\pi\delta^2)^{3/2} \\
&\times \int_{m_e}^{Q_{\beta^+} - m_e} (2\pi) \delta(Q_{\beta^+} - m_e - E_+ - E_\nu) E_\nu \\
&\times \left\{ 1 + \sum_{i>j} 2U_{ei}^* U_{ej} e^{-\delta^2(\Delta\vec{k}_{ij})^2} \right. \\
&\times \cos \left[\left(\sqrt{E_\nu^2 + m_i^2} - \sqrt{E_\nu^2 + m_j^2} \right) t \right] \left. \right\} \\
&\times F(Z-1, E_+) \sqrt{E_+^2 - m_e^2} E_+ dE_+, \quad (14)
\end{aligned}$$

where $\Delta\vec{k}_{ij} = (\vec{k}_i - \vec{k}_j)/2$ and $e^{-\delta^2(\Delta\vec{k}_{ij})^2}$ are kept as input parameters [4].

For the calculation of the r.h.s of Eq.(14) we have set neutrino masses zero everywhere except in the energy difference $E_i(\vec{k}_i) - E_j(\vec{k}_j)$ [4]. Then, due to the exponential function $e^{-\delta^2(\frac{\vec{k}_i - \vec{k}_j}{2})^2}$ the neutrino momenta are constrained by $\vec{k}_i \simeq \vec{k}_j = \vec{k}$ [4]. In such an approximation we get $E_j(\vec{k}_j) \simeq E_\nu = |\vec{k}|$ and $E_i(\vec{k}_i) - E_j(\vec{k}_j) = \sqrt{E_\nu^2 + m_i^2} - \sqrt{E_\nu^2 + m_j^2}$.

In terms of the rate of the neutrino spectrum the time-dependent β^+ -decay rate is defined by [4]

$$\lambda_{\beta^+}(t) = \int \frac{d^3 k}{(2\pi)^3 2E_\nu} \frac{1}{(\pi\delta^2)^{3/2}} \frac{dN_\nu(t)}{dt}. \quad (15)$$




FIG. 2: Time-dependence of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion on the time-interval equal to $2T_{EC} = 14$ s.

For $\theta_{13} = 0$ [7] (see also [4]) we deal with two-neutrino mass-eigenstates and obtain

$$\frac{\lambda_{\beta^+}(t)}{\lambda_{\beta^+}} = 1 + R_{\beta^+}(t), \quad (16)$$

where λ_{β^+} is given by Eq.(4) and $R_{\beta^+}(t)$ is equal to

$$\begin{aligned} R_{\beta^+}(t) = & \sin 2\theta_{12} e^{-\delta^2(\Delta\vec{k}_{12})^2} \int_{m_e}^{Q_{\beta^+}-m_e} dE_+ E_+ \\ & \times \sqrt{E_+^2 - m_e^2} (Q_{\beta^+} - m_e - E_+)^2 \frac{F(Z-1, E_+)}{f(Q_{\beta^+}, Z-1)} \\ & \times \cos \left[\left(\sqrt{(Q_{\beta^+} - m_e - E_+)^2 + m_2^2} \right. \right. \\ & \left. \left. - \sqrt{(Q_{\beta^+} - m_e - E_+)^2 + m_1^2} \right) t \right]. \end{aligned} \quad (17)$$

For the numerical calculations we use $\sin 2\theta_{12} e^{-\delta^2(\Delta\vec{k}_{12})^2} = 0.20$ [4] and $Q_{\beta^+} = 3396(6)$ keV [9].

The time-dependent part of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion on the time-interval equal to $2T_{EC} = 14$ s, i.e. two periods of the time-modulation of the EC -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion, is shown in Fig. 2. For the calculation of $R_{\beta^+}(t)$ we have used neutrino masses $m_j(R)$, obtained in [5] and corrected by the interaction of massive neutrinos with the strong Coulomb field of the daughter nucleus $^{140}\text{Ce}^{58+}$ [5]. It is seen that the β^+ -decay rate varies in time much faster than the EC -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion.

In the measurement of the time-dependence of the β^+ -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ions the time-spectrum of the decay is defined by the time-intervals $\Delta T = 5 \times \Delta T_{\text{bin}}$, caused by 5 bins with the length $\Delta T_{\text{bin}} = 64$ ms each. This leads to the experimental value of the β^+ -decay rate, averaged over the time-interval $\Delta T = 5 \times \Delta T_{\text{bin}} = 0.32$ s. Due to the rapid variations of $R_{\beta^+}(t)$ a modulation of the time-dependence of the β^+ -decay rate $\lambda_{\beta^+}(t)$ of the H-like $^{140}\text{Pr}^{58+}$ ion is not observable in an experiment.

CONCLUSION

We have studied the time-dependence of the β^+ -decay rate $\lambda_{\beta^+}(t)$ of the H-like $^{140}\text{Pr}^{58+}$ ion. For the calculation of $\lambda_{\beta^+}(t)$ we have followed the approach, proposed in [4]–[6] as applied to the explanation of the time-modulation of the EC -decay rate of the H-like $^{140}\text{Pr}^{58+}$ ion. We have found that the time-dependent term of the β^+ -decay rate varies very rapidly in time, which makes such a time-dependence unobservable. This result can be used as a prediction for future analysis of the time-dependence of the β^+ -decay rates of the H-like heavy ions $^{140}\text{Pr}^{58+}$ and $^{142}\text{Pm}^{60+}$ at GSI for the test of the measuring method.

* Electronic address: ivanov@kph.tuwien.ac.at

- [1] Yu. A. Litvinov *et al.* (the GSI Collaboration), Phys. Lett. B **664**, 162 (2008).
- [2] F. Bosch, “Observation of non-exponential Decays of Hydrogen-like ^{140}Pr and ^{142}Pm Ions”, Invited talk at Annual Meeting of ENTApP N6/WP1 “Physics of Massive Neutrinos”, 19 May – 22 May of 2008 on Milos Island, Greece.
- [3] P. Kienle, “Time Modulation of the Electron Capture Decay due to Neutrino Mixing”, Invited talk at Annual Meeting of ENTApP N6/WP1 “Physics of Massive Neutrinos”, 19 May – 22 May of 2008 on Milos Island, Greece.
- [4] A. N. Ivanov, R. Reda, and P. Kienle, *On the time-modulation of the K-shell electron capture decay of H-like $^{140}\text{Pr}^{58+}$ ions produced by neutrino-flavour mixing*, nucl-th/0801.2121.
- [5] A. N. Ivanov, E. L. Kryshen, M. Pitschmann, and P. Kienle, *Neutrino masses from the Darmstadt oscillations*, nucl-th/0804.1311.
- [6] A. N. Ivanov, P. Kienle, E. L. Kryshen, M. Pitschmann, *Massive neutrinos and “Darmstadt oscillations”*, Invited talk at Annual Meeting of ENTApP N6/WP1 “Physics of Massive Neutrinos”, 19 May – 22 May of 2008 on Milos Island, Greece.
- [7] W.-M. Yao *et al.*, J. Phys. G **33**, 156, 471 (2006).
- [8] Yu. A. Litvinov *et al.* (the GSI Collaboration) Phys. Rev. Lett. **99**, 262501 (2008).
- [9] A. N. Ivanov, M. Faber, R. Reda, and P. Kienle, *Weak decays of H-like $^{140}\text{Pr}^{58+}$ and He-like $^{140}\text{Pr}^{57+}$ ions*, nucl-th/0711.3184 (to appear in PRC).
- [10] N. Cottingham and D. A. Greenwood, in *An introduction to nuclear physics*, Cambridge University Press, Second Edition, 2001.
- [11] H. F. Schopper, in *Weak interactions and nuclear beta decay*, North-Holland Publishing Co., Amsterdam, 1966.